

An Ounce of Prevention Emergency Warning Signs

In adults, emergency warning signs that need urgent medical attention include:

- ◆ Difficulty breathing or shortness of breath
- ◆ Pain or pressure in the chest or abdomen
- ◆ Sudden dizziness
- ◆ Confusion
- ◆ Severe or persistent vomiting
- ◆ Flu-like symptoms improve but then return with fever and worse cough

In children, emergency warning signs that need urgent medical attention include:

- ◆ Fast breathing or trouble breathing
- ◆ Bluish or gray skin color
- ◆ Not drinking enough fluids
- ◆ Severe or persistent vomiting
- ◆ Not waking up or not interacting
- ◆ Being so irritable that the child does not want to be held
- ◆ Flu-like symptoms improve but then return with fever and worse cough



**Working
together
to keep
our seniors
healthy**

America's Choice in Homecare®



- Up to 24 Hour Care
- Meal Preparation
- Hygiene Assistance
- Light Housekeeping
- Errands
- Shopping
- Doctor Appointments
- Respite for Family Caregivers
- Rewarding Companionship

Solution...

VisitingAngels®
LIVING ASSISTANCE SERVICES
Select Your Caregiver!



**An Ounce of
Prevention
H1N1**



**Prevention and
Treatment**



**A guide for family, friends
and loved ones**
information provided by



from
America's Choice in Homecare®



For more information visit: www.cdc.gov/h1n1flu

www.visitingangels.com

A Public Service from Visiting Angels®

Each agency privately owned and operated.

H1N1: Prevention & Treatment

A complete guide on the prevention and treatment of the H1N1 flu from the Centers for Disease Control

What can I do to protect myself from getting sick?

There is no vaccine available right now to protect against 2009 H1N1 virus. However, a 2009 H1N1 vaccine is currently in production and may be ready for the public in the fall. As always, a vaccine will be available to protect against seasonal influenza. There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza.

Take these everyday steps to protect your health:

- ◆ Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- ◆ Wash your hands often with soap and water. If soap and water are not available, use an alcohol based hand rub - sanitizer.
- ◆ Avoid touching your eyes, nose or mouth. Germs spread this way.
- ◆ Try to avoid close contact with sick people.
- ◆ If you are sick with flu-like illness, the CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from others as much as possible to keep from making others sick.

Other important actions that you can take are:

- ◆ Follow public health advice regarding school closures, avoiding crowds and other social distancing measures.
- ◆ Be prepared in case you get sick and need to stay home for a week or so; a supply of over-the-counter medicines, alcohol-based hand rubs (for when soap and water are not available), tissues and other related items could help you to avoid the need to make trips out in public while you are sick and contagious.

If I have a family member at home who is sick with 2009 H1N1 flu, should I go to work?

Employees who are well but who have an ill family member at home with 2009 H1N1 flu can go to work as usual. These employees should monitor their health every day, and take everyday precautions including washing their hands often with soap and water, especially after they cough or sneeze. If soap & water are not available, they should use an alcohol-based hand rub. If they become ill, they should notify their supervisor and stay home. Employees who have an underlying medical condition or who are pregnant should call their health care provider for advice, because they might need to receive influenza antiviral drugs to prevent illness. For more information please see General Business and Workplace Guidance for the Prevention of Novel Influenza A (H1N1) Flu in Workers.

What is the best technique for washing my hands to avoid getting the flu?

Washing your hands often will help protect you from germs. CDC recommends that when you wash your hands -- with soap and warm water -- that you wash for 15 to 20 seconds. When soap and water are not available, alcohol-based disposable hand wipes or gel sanitizers may be used. You can find them in most supermarkets and drugstores. If using gel, rub your hands until the gel is dry. The gel doesn't need water to work; the alcohol in it kills the germs on your hands.

Contamination - Cleaning

What household cleaning should be done to prevent the spread of influenza virus?

To prevent the spread of influenza virus it is important to keep surfaces (especially bedside tables, surfaces in the bathroom, kitchen counters and toys for children) clean by wiping them down with a household disinfectant according to directions on the product label.

All content provided within this informational brochure is from the Centers for Disease Control and Prevention.
Visit www.cdc.gov/h1n1flu for more information

What should I do if I get sick?

If you live in areas where people have been identified with 2009 H1N1 flu and become ill with influenza-like symptoms, including fever, body aches, runny or stuffy nose, sore throat, nausea, or vomiting or diarrhea, you should stay home and avoid contact with other people. CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Stay away from others as much as possible to keep from making others sick. Staying at home means that you should not leave your home except to seek medical care. This means avoiding normal activities, including work, school, travel, shopping, social events, and public gatherings.

Are there medicines to treat 2009 H1N1 infection?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with 2009 H1N1 flu virus. Antiviral drugs are prescription medicines (pills, liquid or an inhaled powder) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. During the current pandemic, the priority use for influenza antiviral drugs is to treat severe influenza illness (for example hospitalized patients) and people who are sick who have a condition that places them at high risk for serious flu-related complications.

